

Antiglycation and Antioxidant Activities of Black Garlics in Streptozotocin and Staphylococcus aureus Induced Rats Model of Diabetes Mellitus

by Evy Yulianti, Kartika Ratna Pertiwi, Tutiek Rahayu

ABSTRACT

Diabetes mellitus (DM) is one of the diseases that has received attention due to various complications and is one of the comorbidities of the severity of Covid-19. Free radicals are products of oxidative stress that arise because of hyperglycemia, so the prevention of their formation and their elimination is a potential mechanism in the management of DM patients. One biological potential candidate reported to have anti-hyperglycemic effects is *black garlic*. This study aims to determine the antiglycation and antioxidant activity of black garlics in DM model rats induced with *streptozotocin* (STZ) and infected with *S aureus*. The study used male Wistar rats (8 weeks, 200 g) which were divided into 7 groups, namely normal mice, STZ-induced mice, STZ and *S aureus*-induced rat and 3 groups of STZ and *S aureus*-induced rats treated with 3 doses of black garlics and 1 metformin dose. The parameters measured in this study were glucose levels measured using the GOD PAP method; H₂O₂ with spectrophotometry, and CML with ELISA. The results showed that black garlic ethanol extract has the potential as an antiglycation agent in terms of the IC value of ₅₀ BSA-glucose test. The administration of black garlics ethanol extract for 7 days at levels of 1.5, 3 and 6 mg / 200gBW can reduce plasma glucose levels and levels of 6 mg / 200gBW can reduce H₂O₂.

Kata Kunci: *diabetes, streptozotocin, S aureus, black garlic, antiglycation, antioxidant*